

Central Exclusive Meson Pair Production in the Perturbative Regime

Lucian Harland-Lang


Cavendish Laboratory, University of Cambridge

DIS 2011, Newport News, VA USA, April 11-15, 2011.

Based on work by V.A. Khoze, M.G. Ryskin, W.J. Stirling and L.A. Harland-Lang. (KHYSTHAL collaboration)
(Paper in preparation)

Central Exclusive Production (CEP) : Durham model

- Colliding protons interact via a colour singlet exchange and remain intact: can be measured by adding detectors far down the beam-pipe.
- A system X of mass M_X is produced at the collision point, and *only* its decay products are present in the central detector.
- The generic process $pp \rightarrow p + X + p$ is modeled perturbatively by the exchange of two t-channel gluons, with the use of pQCD justified by the presence of a hard scale $\sim M_X$.
- ‘ $J_z = 0$ selection rule’: production of states with non- $J_z^P = 0^+$ quantum numbers is strongly suppressed by ~ 2 orders of magnitude.



Meson pair CEP

- CEP is a promising way to study new physics at the LHC, but we can also consider the CEP of lighter, established objects : χ_c , $\gamma\gamma$ and jj CEP already observed at the Tevatron.
- Can serve as 'Standard Candle' processes, which allow us to check the theoretical predictions for central exclusive new physics signals at the LHC, as well as being of interest in their own right¹.
- This talk will focus on the CEP of light ($q = (u, d, s)$, in $m_q = 0$ limit) meson pairs, $M\bar{M}$, at sufficiently high invariant mass for perturbative formalism to be applicable:
 - ▶ Provides novel application/test of hard exclusive formalism, complementary to more standard photon-induced processes ($\gamma\gamma \rightarrow M\bar{M}$, $\gamma\gamma^{(*)} \rightarrow M$ etc²).
 - ▶ Demonstrates application of MHV formalism to simplify/check calculations.
 - ▶ $\pi^0\pi^0$ CEP a possible background to $\gamma\gamma$ CEP.
 - ▶ Could probe the $q\bar{q}$ and $g\bar{g}$ content of η , η' mesons?
 - ▶ An interesting potential observable @ RHIC, Tevatron and LHC: meson pair CEP data (at lower p_\perp) already being taken by ALICE and CDF.

¹ See LHL, V.A. Khoze, M.G. Ryskin, W.J. Stirling, [arXiv:1005.0695](#) and [arXiv:1011.0680](#).

² For a recent review, see for example V. L. Chernyak, [arXiv:0912.0623](#).



Modeling meson pair CEP perturbatively

- Simpler exclusive process $\gamma\gamma \rightarrow M\bar{M}$ ($= \pi^0\pi^0, \pi^+\pi^-, K^+K^- \dots$) at large angles was calculated ~ 30 years ago³.
- Total amplitude given by convolution of parton level $\gamma(\lambda_1)\gamma(\lambda_2) \rightarrow q\bar{q}q\bar{q}$ amplitude with non-perturbative pion wavefunction $\phi(x)$

$$\mathcal{M}_{\lambda_1\lambda_2}(s, t) = \int_0^1 dx dy \phi(x)\phi(y) T_{\lambda_1\lambda_2}(x, y; s, t)$$

where helicity amplitudes $T_{\lambda_1\lambda_2}$ can be calculated perturbatively.

- With suitable choice of $\phi(x)$ shape, $\gamma\gamma \rightarrow M\bar{M}$ data are described quite well (see plot⁴).



³S. J. Brodsky and G. P. Lepage, Phys. Rev. D 24 (1981) 1808.

⁴Data taken from Belle Collaboration, Phys. Lett. B615 (2005) 39

$gg \rightarrow M\bar{M}$ calculation

- Simplest case: production of flavour non-singlet scalar mesons (e.g. $\pi^0\pi^0, \pi^+\pi^-\dots$).
- Can calculate the LO $gg \rightarrow M\bar{M}(= q\bar{q}q\bar{q})$ amplitudes to give

$$T_{++} = T_{--} = 0 ,$$
$$T_{-+} = T_{+-} \propto \frac{\alpha_S^2}{a^2 - b^2 \cos^2 \theta} \left(\frac{N_c}{2} \cos^2 \theta - C_F a \right) ,$$

where $a, b = (1 - x)(1 - y) \pm xy$.

- $J_Z = 0$ amplitudes vanish, as in $\gamma\gamma \rightarrow M\bar{M}$ for neutral mesons. We therefore expect a strong suppression in flavour non-singlet $M\bar{M}$ CEP due to $J_Z = 0$ selection rule.
- $J_Z = 2$ amplitudes contain 'radiation zero', vanishing for a physical value of $\cos^2 \theta$. Well known effect in all gauge theories (e.g. $u\bar{d} \rightarrow W^+\gamma$), but usually washed out in QCD by colour averaging. Here, position of zero depends on the choice of $\phi(x)$, and we find that there is always a zero in the physical region for any choice of $\phi(x)$ and general N_c .

$gg \rightarrow M\bar{M}$ amplitude: Feynman diagrams

Vanishing of T_{++} , T_{--} follows after calculating:

$$\frac{1}{2} C_F \frac{\delta^{AB}}{N_c} \left(\begin{array}{c} \text{Diagram 1} \\ \text{Diagram 2} \\ \text{Diagram 3} \\ \text{Diagram 4} \end{array} \right) + \frac{1}{2} C_F \frac{\delta^{AB}}{N_c} \left(\begin{array}{c} \text{Diagram 5} \\ \text{Diagram 6} \\ \text{Diagram 7} \\ \text{Diagram 8} \end{array} \right) - \frac{N_c}{4} \frac{\delta^{AB}}{N_c} \left(\begin{array}{c} \text{Diagram 9} \\ \text{Diagram 10} \\ \text{Diagram 11} \end{array} \right)$$

$gg \rightarrow M\bar{M}$ amplitude: MHV calculation (1)

- $g(+)\bar{g}(+) \rightarrow q(\pm)\bar{q}(\mp)q(\pm)\bar{q}(\mp)$ amplitude is MHV: maximum $(n-2)$ number of particles have same helicity.
- Such amplitudes known to have remarkably simple forms, and corresponding 'spinor helicity' formalism can greatly simplify calculation.
- T_{++}, T_{--} can be calculated from known Parke-Taylor amplitude⁵

$$M_n \propto \sum_{\sigma} \frac{\langle k_p k_{\bar{q}} \rangle}{\langle k_p a_1 \rangle \cdots \langle a_l k_{\bar{q}} \rangle} \frac{\langle k_q k_{\bar{p}} \rangle}{\langle k_q b_1 \rangle \cdots \langle b_{l'} k_{\bar{p}} \rangle} (\lambda^{a_1} \cdots \lambda^{a_l})_{i_1 j_2} (\lambda^{b_1} \cdots \lambda^{b_{l'}})_{i_2 j_1} \\ - \frac{1}{N_c} \frac{\langle k_p k_{\bar{p}} \rangle}{\langle k_p a_1 \rangle \cdots \langle a_l k_{\bar{p}} \rangle} \frac{\langle k_q k_{\bar{q}} \rangle}{\langle k_q b_1 \rangle \cdots \langle b_{l'} k_{\bar{q}} \rangle} (\lambda^{a_1} \cdots \lambda^{a_l})_{i_1 j_1} (\lambda^{b_1} \cdots \lambda^{b_{l'}})_{i_2 j_2} .$$

- Making colour singlet identification ($i_1 = j_2, i_2 = j_1$) and identifying $q\bar{q}, p\bar{p}$ with collinear quarks within mesons

$$k_q = xk_3 \quad k_{\bar{q}} = (1-y)k_4 \quad k_p = yk_4 \quad k_{\bar{p}} = (1-x)k_3 ,$$

then amplitude reduces to


$$M \propto \langle k_3 k_2 \rangle \langle k_1 k_4 \rangle + \langle k_1 k_3 \rangle \langle k_2 k_4 \rangle - \langle k_3 k_4 \rangle \langle k_1 k_2 \rangle = 0 ,$$

which vanishes from the Schouten identity.

⁵M. L. Mangano, S. J. Parke, Phys. Rept. 200 (1991) 301-367

$gg \rightarrow M\bar{M}$ amplitude: MHV calculation (2)


- The vanishing of the $gg \rightarrow M\bar{M}$ $J_Z = 0$ amplitudes follow directly from the corresponding 6-particle MHV amplitude. This result depends crucially on the colour singlet projection and collinearity of the $q\bar{q}$ pairs, and only occurs for non-flavour singlet mesons
- The MHV formalism can be extended to include the non-MHV $|J_Z| = 2$ amplitude: contributing amplitudes given by tree graphs in which the vertices are the usual tree-level MHV scattering amplitudes continued off-shell⁶.
- More complicated than $J_Z = 0$ case, but an explicit calculation within this framework confirms our result.



⁶see e.g. G. Georgiou, V. V. Khoze, JHEP 0405 (2004) 070. [[hep-th/0404072](#)].

Flavour singlet meson production


- A second set of diagrams can in general contribute, where the $q\bar{q}$ forming the mesons connected by a quark line (no equivalent diagram in $\gamma\gamma \rightarrow M\bar{M}$ process).
- Only relevant for flavour singlet states (e.g. for $gg \rightarrow \pi^0\pi^0$, $|u\bar{u}\rangle$ and $|d\bar{d}\rangle$ Fock components interfere destructively).
- In this case the $J_z = 0$ amplitude does not vanish \rightarrow Expect strong enhancement in $\eta'\eta'$ CEP rate⁷ and (through η - η' mixing), some enhancement to $\eta\eta$ rate. $\eta\eta'$ CEP can also occur via this mechanism.
- Also: any sizable gg component to flavour singlet states, contributing through $gg \rightarrow 4g$ and $gg \rightarrow q\bar{q}gg$ processes, may in principle strongly enhance the CEP cross section (again $J_z = 0$ amplitudes do not vanish). A significant 'excess' in future CEP data could be evidence for this.



⁷Recall quark content of $|\eta'\rangle$ is dominantly $\sim |u\bar{u} + d\bar{d} + s\bar{s}\rangle$



Numerical Results

- Strong enhancement in flavour singlet states clear, with precise η'/η hierarchy given by choice of $\eta - \eta'$ mixing angle.
- CEP cross sections for vector mesons ($\rho\rho$, $\omega\omega$, $\phi\phi$) can be calculated in the same way.



- $\pi^0\pi^0$ CEP can in principle be an important background to $\gamma\gamma$ CEP, but we find this not to be the case. This depends crucially on vanishing of the $gg \rightarrow \pi^0\pi^0$ amplitude for $J_Z = 0$ initial-state gluons.
- However: possible $J_Z = 0$ contribution from higher twist effects, NNLO corrections... could increase flavour non-singlet rate by a factor 'a few'. Also, possible non-perturbative contribution at lower p_\perp ?

\overline{MM} CEP: secondary mechanism



- As well as the standard CEP diagram (a), we must in principle consider the process shown in diagram (b)⁸.
- It is formally **subleading**, as the amplitude has an extra power of the pion transverse momentum squared, k_{\perp}^2 , in the denominator.
- However: we have seen that CEP $\pi^0\pi^0$ process is strongly suppressed by the $J_Z = 0$ selection rule, so this diagram may be important.
- Preliminary numerical estimates suggest this *may* be the case at lower p_{\perp} .

⁸ackn. to Jeff Forshaw for pointing this out.

A MC event generator including⁹:

- Simulation of different CEP processes, including all spin correlations:
 - $\chi_{c(0,1,2)}$ CEP via the $\chi_c \rightarrow J/\psi \gamma \rightarrow \mu^+ \mu^- \gamma$ decay chain.
 - $\chi_{b(0,1,2)}$ CEP via the equivalent $\chi_b \rightarrow \Upsilon \gamma \rightarrow \mu^+ \mu^- \gamma$ decay chain.
 - $\chi_{(b,c)J}$ and $\eta_{(b,c)}$ CEP via general two body decay channels
 - Physical proton kinematics + survival effects for quarkonium CEP at RHIC.
 - $\gamma\gamma$ CEP.
 - Exclusive J/ψ and Υ photoproduction.
 - **Meson pair CEP** ($\pi^0\pi^0$, $\eta\eta$, $\eta'\eta'$...) to be included soon.
 - More to come (dijets, open quark, Higgs...?).
- Via close collaboration with CDF, STAR and LHC groups, in both proposals for new measurements and applications of SuperCHIC, it is becoming an important tool for current and future CEP studies.


⁹The SuperCHIC code and documentation are available at
<http://projects.hepforge.org/superchic/>

Summary and Outlook

- CEP in hadron collisions offers a promising framework within which to study novel aspects of QCD and new physics signals.
- CEP processes observed at the Tevatron, RHIC and early LHC can serve as ‘standard candles’ for new physics CEP at the LHC.
- The CEP of mesons pairs at high invariant masses is an interesting process, representing a novel application of pQCD framework for describing exclusive processes (publication in preparation).
- MHV formalism can be readily applied to these processes, and can simplify calculations.
- CEP could help probe the gluonic structure of η , η' mesons.
- Perturbative calculation predicts that $\pi^0\pi^0$ BG to $\gamma\gamma$ CEP is suppressed.
- More CEP results to come from RHIC, the Tevatron and LHC in the future.

Backup 1: radiation zeros


- Complete destructive interference of radiation patterns, resulting in vanishing amplitude for certain configuration of final state particles.
- Occurs in most Born amplitudes for radiation of massless gauge bosons, first seen in $u\bar{d} \rightarrow W^+\gamma$ amplitude.
- General conditions for zeros are known¹⁰: often zeros do not occur in physical region.
- Occurs in QCD, but zeros are usually neutralised along with colour by averaging of hadronisation \rightarrow pure colour singlet CEP process *in principle* uniquely positioned to observe zeros.
- **However**: zero only occurs at LO in subleading $|J_Z| = 2$ amplitude. We may reasonably expect higher order ($J_Z = 0$) corrections to fill in the zero.



¹⁰S.J. Brodsky and R.W. Brown, Phys. Rev. Lett. 49, 966 (1982)

Backup 2: $M\bar{M}$ non-perturbative production

- For low values of pion p_{\perp} , expect non-perturbative double Pomeron/Reggeon exchange mechanism to contribute, mediated via an off-shell pion.



- Uncertainty in what to take for form factor of off-shell pion ('soft' vs 'hard' fit), which suppresses high values of final state pion p_{\perp} , leads to quite large uncertainty in expected rate.
- Expect smooth transition with increasing p_{\perp} between non-perturbative (dominantly real amplitude) and perturbative (dominantly imaginary amplitude) dominance.
- Measurement of $\pi^0\pi^0/\pi^+\pi^-$ CEP in low p_{\perp} region would help constrain off-shell pion form factor.